CLAIMS

We Claim:

10

15

- 5 1. An isolated polymer comprising chitosan bound to a glutamine residue of a polypeptide.
 - 2. An isolated polymer comprising chitosan bound to a tyrosine residue of a polypeptide.
 - 3. The polymer of claim 1 or 2, wherein the polypeptide is not a polypeptide that is covalently bonded to chitosan in surimi.
 - 4. The polymer of claim 1 or 2, wherein the polypeptide is gelatin.
 - 5. The polymer of claim 1 or 2, wherein the polypeptide is 2-5 amino acids in length.
- 6. The polymer of claim 1 or 2, wherein the polypeptide is 6-20 amino acids in length.
 - 7. The polymer of claim 1 or 2, wherein the polypeptide is at least 21 amino acids in length.
- 8. The polymer of claim 1 or 2 wherein the polymer is chitosan covalently bonded to one or more polypeptides, a polypeptide covalently bonded to one or more chitosan, or a combination thereof.
- 9. The polymer of claim 8 wherein the chitosan is bound to two or more30 polypeptides of different types.
 - 10. A composition comprising a polysaccharide, a polypeptide and an enzyme, wherein the enzyme is transglutaminase or tyrosinase.

11. A method of making a polymer, which comprises contacting a polypeptide and a polysaccharide with a transglutaminase under conditions sufficient to bind the polypeptide to the polysaccharide, wherein the polypeptide comprises a glutamine residue.

5

12. A method of making a polymer, which comprises:

contacting a polypeptide comprising a glutamine residue with transglutaminase to yield a modified polypeptide; and

contacting the modified polypeptide with a polysaccharide under conditions sufficient to bond the polysaccharide to the modified polypeptide.

13. A method of making a polymer, which comprises contacting a polypeptide and a polysaccharide with a tyrosinase under conditions sufficient to bind the polypeptide to the polysaccharide, wherein the polypeptide comprises a tyrosine residue.

15

20

14. A method of making a polymer, which comprises:

contacting a polypeptide comprising a tyrosine residue with tyrosinase to yield a modified polypeptide; and

contacting the modified polypeptide with a polysaccharide under conditions sufficient to bond the polysaccharide to the modified polypeptide.

- 15. The method of one of claims 10-14, wherein the polypeptide is gelatin.
- 16. The method of one of claims 10-14, wherein the polysaccharide is chitosan.